

Risk theory and risk management in actuarial science

Winter term 2017/18

2nd work sheet

13. Consider a portfolio consisting of the following index futures *Dow Jones Industrial Average* (\hat{DJI}), *S&P 500 ETF* (\hat{GSPC}), *Nasdaq Composite* (\hat{IXIC}), *DAX* (\hat{GDAXI}) and *ATX* (\hat{ATX}) with one piece per each index future. Estimate the weekly value at risk $\text{VaR}_{0.90}$ of this portfolio in two ways as described below. Then compare and comment on the obtained results.
- (a) Use historical simulation of the losses over the last 10 years, from October 20, 2007 until October 20, 2017.
 - (b) Use the variance-covariance method, again based on the weekly logarithmic returns of the last 10 years as in (a).

The data can be downloaded from finance.yahoo.com: search for the required index future (you can well search for the abbreviations given in paranthesis above), click 'Historical Prices', update the 'Time Period' and 'Frequency' appropriately, and finally click on 'Download data'. Use the adjusted close prices to compute the weakly logarithmic returns of the single index futures.

14. Use the bootstrapping method to compute a centered confidence interval with confidence level 10% for the $\text{VaR}_{0.95}$ and $\text{CVaR}_{0.95}$ of the weakly loss of a portfolio given as described in Exercise 13. Experiment with different values of the parameter N which specifies the number of repetitions of bootstrapping while calculating one estimator per repetition.
15. Consider the daily logarithmic returns (based on adjusted close prices) of the BMW and Siemens assets, *BMW.F* and *SIA.AS*, respectively, over the time interval October 20, 2007, and October 20, 2017. Use the Hill estimator to get an approximation of the coefficient of the regular variation and determine the corresponding estimates for $\text{VaR}_{0.90}$ and $\text{CVaR}_{0.90}$ for each of these assets. Specify a plausible range of values of k to be chosen (depending on the sample size) and generate the Hill plot for those values of k . Based on the Hill plot make a suggestion for an appropriate value of k to be used and argue your choice carefully. Use yahoo.finance.com as a data source (see Exercise 13).
16. By means of the qq-plot check whether a normal distribution or a heavy tailed distribution like the (generalized) Pareto distribution is more appropriate to model the right tail of the loss distribution of the BMW and Siemens assets as described in Exercise 15, respectively. To this end you should compare the empirical quantiles of the above mentioned losses to the (numerically or analytically) computed quantiles of the reference distributions (i.e. a normal and a generalized Pareto distribution) and summarize the results graphically as described schematically in the lecture.